

SUPERSYMMETRIC HIGGS BOSONS IN A 5D ORBIFOLD MODEL

V. DI CLEMENTE AND S. F. KING

*Department of Physics and Astronomy,
University of Southampton, Southampton, SO17 1BJ, U.K.
E-mail: vicente@hep.phys.soton.ac.uk, sfk@hep.phys.soton.ac.uk*

D. A. J. RAYNER

*Dipartimento di Fisica ‘G. Galilei’, Università di Padova and INFN,
Sezione di Padova, Via Marzola 8, I-35131 Padua, Italy
E-mail: rayner@pd.infn.it*

We analyze the phenomenology of the Higgs sector in a 5D model compactified on an S_1/Z_2 orbifold with a compactification scale $M_C \sim \mathcal{O}(\text{TeV})$ where supersymmetry breaking is localized on a brane at one of the fixed points. We show that the conventional MSSM Higgs boson mass bounds in 4D can be violated when we allow the gauge sector, Higgs and third family multiplets to live in the fifth extra dimension.

Introduction

Extra-dimensional supersymmetric models with a TeV compactification scale M_C offer an exciting new environment for investigating electroweak symmetry breaking (EWSB) ¹. Higher-dimensional fields can be equivalently described as infinite towers of Kaluza-Klein (KK) resonances and the k^{th} KK mode has a mass $m_k \sim \mathcal{O}(k/R)$, where $R = M_C^{-1}$ is the compactification radius of the extra dimension(s) ^a. These models have the attractive feature that loops of virtual top and stop KK modes yield a negative contribution to the Higgs mass, which can trigger EWSB around the TeV scale. Remarkably, the one-loop effective potential is found to be finite due to $\mathcal{N} = 1$ SUSY in 5D which implies that the Higgs physics is completely independent of the high-energy physics above some cutoff scale.

^aNote that $\mathcal{N} = 1$ SUSY in 5D is equivalent to $\mathcal{N} = 2$ SUSY in 4D, and so 5D hypermultiplets can be decomposed into an $\mathcal{N} = 1$ multiplet plus its mirror-conjugate.

Our Model

We will summarize the features of our 5D model ² as shown in Figure 1. We localize 4D 3-branes at the two fixed points $y = 0, \pi R$ arising from the compactification of the extra dimension on an S^1/Z_2 orbifold. SUSY is broken on the “source” brane at $y = \pi R$ when a localized gauge field singlet ($\hat{S}$) acquires a non-zero F-term vacuum expectation value (VEV).




Figure 1. The allocation of states in our 5D model is motivated by an explicit type I string construction ³.

The first two MSSM ($\hat{F}_{1,2}$) families are confined to the “Yukawa” brane at $y = 0$ where we are forced to localize Yukawa couplings, while the third family ($\hat{F}_3$), MSSM gauge sector ($\hat{\mathcal{V}}_{MSSM}$) and Higgs superfields ($\hat{H}_{u,d}$) live in the extra dimensional bulk along with their mirror-conjugate partners. Bulk fields acquire tree-level soft parameters due to their direct coupling with the SUSY breaking brane through non-renormalizable operators defined at $y = \pi R$, while first and second family squarks are suppressed by the separation between branes ^{4,5} which alleviates the flavour-changing neutral current problem. The electroweak symmetry is broken when the zero-modes of the Higgs fields acquire non-vanishing VEVs ($\langle h_u \rangle, \langle h_d \rangle \neq 0$).

The localization of the Yukawa couplings and soft SUSY breaking masses at the fixed points leads to complicated mixing between different KK modes when we decompose the 5D fields ^b. In Ref. ², we diagonalized the top/stop mass matrices to find the field-dependent KK mass spectra

^bThe mixing strength is related to the SUSY breaking VEVs, and viable radiative EWSB requires a non-universality between VEVs, i.e. $F_{S,\tilde{t}} \gg F_{S,H}$.

for the top:

$$m_{t,k}[h_u] = \left| k M_C + \frac{M_C}{\pi} \arctan \left(\frac{y_t h_u \pi}{\sqrt{2} M_C} \right) \right| \quad (k = -\infty, \dots, \infty) \quad (1)$$

and the field-independent stop KK modes:

$$m_{\tilde{t},k}^2 \approx \left(k + \frac{1}{2} \right)^2 M_C^2 \quad (k = -\infty, \dots, \infty) \quad (2)$$

where y_t is the 4D Yukawa coupling and we identify the physical top mass as the $k = 0$ mode of Eq. 1. We used dimensional regularization and zeta-function regularization techniques⁶ to evaluate the top/stop contributions to the 1-loop effective potential, and found that the top and stop contributions are separately finite (and free of ultraviolet divergences) due to supersymmetric cancellations.

Results

We applied the usual minimization conditions to the 1-loop effective potential to extract the physical Higgs mass spectrum in terms of a universal soft Higgs mass input parameter (or equivalently m_{A^0}) and $\tan \beta$. The results are shown in Figure 2 for two different values of $\tan \beta$ where fine-tuning arguments⁷ restrict our analysis to compactification scales $M_C \lesssim 4$ TeV. We also compare the MSSM results⁸ with our model for $\tan \beta = 1.5$ and find that our model is *not* excluded by the LEP signal⁹:




Figure 2. The Higgs masses ($m_{h^0}, m_{H^\pm}, m_{H^0}$) against m_{A^0} for $\tan \beta = 1.5$ (left panel) and $\tan \beta = 20$ (right panel). The LEP candidate at 115 GeV (*bold dotted-line*) is shown for comparison with m_{h^0} . For $\tan \beta = 1.5$ (left panel), we also compare the MSSM results taken from (*dotted-lines*) against our model (*bold-lines*).




Figure 3. Mass contour plot for the lightest Higgs mass m_{h^0} as a function of the compactification scale M_C (GeV) and $\tan \beta \leq 20$.

In Figure 3, we plot the lightest Higgs mass (m_{h^0}) as a function of the compactification scale M_C and $\tan \beta$. The LEP data excludes the parameter space below the first contour at $m_{h^0} = 115$ GeV⁹ which corresponds to a compactification scale $M_C \approx 1.5 - 1.7$ TeV over the whole range of $\tan \beta$. Notice that our model can easily accommodate the conventional 4D MSSM upper bound on the lightest Higgs boson mass $m_{h^0} \sim 130$ GeV, and can be pushed as high as $m_{h^0} \sim 160$ GeV with $M_C \approx 4$ TeV and $5 \lesssim \tan \beta \lesssim 20$. Remember that including additional matter (e.g. gauge singlets in the NMSSM) can also raise the upper bound on the lightest Higgs mass, but our “minimal” extension of the MSSM in 5D achieves higher mass bounds without adding extra matter content.

References

1. A. Pomarol and M. Quiros, Phys. Lett. B **438** (1998) 255 [arXiv:hep-ph/9806263]; I. Antoniadis, S. Dimopoulos, A. Pomarol and M. Quiros, Nucl. Phys. B **544** (1999) 503 [arXiv:hep-ph/9810410]; A. Delgado, A. Pomarol and M. Quiros, Phys. Rev. D **60** (1999) 095008 [arXiv:hep-ph/9812489]; R. Barbieri, L. J. Hall and Y. Nomura, Phys. Rev. D **63** (2001) 105007 [arXiv:hep-ph/0011311]; N. Arkani-Hamed, L. J. Hall, Y. Nomura, D. R. Smith and N. Weiner, Nucl. Phys. B **605** (2001) 81 [arXiv:hep-ph/0102090]; A. Delgado and M. Quiros, Nucl. Phys. B **607** (2001) 99 [arXiv:hep-ph/0103058].
2. V. Di Clemente, S. F. King and D. A. J. Rayner, Nucl. Phys. B **646** (2002) 24 [arXiv:hep-ph/0205010]; Nucl. Phys. B **617** (2001) 71 [arXiv:hep-ph/0107290].

3. S. F. King and D. A. J. Rayner, Nucl. Phys. B **607** (2001) 77 [arXiv:hep-ph/0012076].
4. D. E. Kaplan, G. D. Kribs and M. Schmaltz, Phys. Rev. D **62** (2000) 035010 [arXiv:hep-ph/9911293]; Z. Chacko, M. A. Luty, A. E. Nelson and E. Ponton, JHEP **0001** (2000) 003 [arXiv:hep-ph/9911323].
5. L. Randall and R. Sundrum, Nucl. Phys. B **557** (1999) 79 [arXiv:hep-th/9810155].
6. V. Di Clemente and Y. A. Kubyshin, Nucl. Phys. B **636** (2002) 115 [arXiv:hep-th/0108117].
7. S. Dimopoulos and G. F. Giudice, Phys. Lett. B **357** (1995) 573 [arXiv:hep-ph/9507282]; G. L. Kane and S. F. King, Phys. Lett. B **451** (1999) 113 [arXiv:hep-ph/9810374], and references therein.
8. D. E. Groom *et al.* [Particle Data Group Collaboration], Eur. Phys. J. C **15** (2000) 1.
9. LEP Higgs Working Group for Higgs boson searches Collaboration, arXiv:hep-ex/0107029.